

Linearization of analytic order relations

Vladimir Kanovei ^{*†‡}

April 1997

Abstract

We prove that if $\preceq$ is an analytic partial order then either $\preceq$ can be extended to a Δ_2^1 linear order similar to an antichain in $2^{<\omega_1}$ ordered lexicographically or a certain Borel partial order $\leq_0$ embeds in $\preceq$. Some corollaries for analytic equivalence relations are given, for instance, if E is a $\Sigma_1^1[z]$ equivalence relation such that E_0 does not embed in E then E is determined by intersections with E -invariant Borel sets coded in $L[z]$.

Introduction.....	2
1. Preliminaries.....	6
2. Monotone Borel functions and the dichotomy.....	8
3. The basic forcing.....	9
4. The product forcing.....	12
5. The construction of an embedding.....	14
6. Why embedding $\leq_0$ is absolute.....	17
7. Borel and analytic order relations.....	18
8. Special cases: Borel classes and generic models.....	21
References.....	24

^{*} Moscow Transport Engineering Institute

[†] kanovei@mech.math.msu.su and kanovei@math.uni-wuppertal.de

[‡] This paper was accomplished in part during my visit to Caltech in April 1997. I thank Caltech for the support and A. S. Kechris and J. Zapletal for useful information and interesting discussions relevant to the topic of this paper during the visit.

Introduction

It is a simple application of Zorn's lemma that any partial order can be extended to a linear order on the same domain. More generally any partial quasi-order admits a linearization.¹

A much more difficult problem is to provide a descriptive characterization of the linear order in the assumption that one has such for the given p. q.-o.. For instance, not every Borel p. q.-o. is *Borel linearizable*.

Example 1. Recall that E_0 is an equivalence relation on 2^ω defined as follows: $a E_0 b$ iff $a(k) = b(k)$ for all but finite k . Let $\preceq$ be E_0 considered as a p. q.-o.. Then $\preceq$ is not Borel linearizable. Indeed any linearization h satisfies $a E_0 b \iff h(a) = h(b)$, but it is known that Borel maps h with such a property do not exist (see Harrington, Kechris, Louveau [2]). $\dashv$

Example 2. Example 1 can be converted to a partial order. Define the *anti-lexicographical* p. o. $\leq_0$ on 2^ω as follows: $a \leq_0 b$ iff either $a = b$ or there is $m \in \omega$ such that $a(k) = b(k)$ for all $k > m$ and $a(m) < b(m)$. Clearly $a \leq_0 b$ implies $a E_0 b$ and $\leq_0$ linearly orders each E_0 -equivalence class similarly to the integers $\mathbb{Z}$, except for the class of $\omega \times \{0\}$ (ordered as ω) and the class of $\omega \times \{1\}$ (ordered as ω^{-1} – the inverted ω)² Finally $\leq_0$ is not Borel linearizable (see Subsection 7.1). $\dashv$

There are Borel-non-linearizable Borel orders of different nature, *e. g.* the p. q.-o. $a \preceq b$ iff $a(k) \leq b(k)$ for all but finite k on 2^ω or the dominance relation on ω^ω . However by the next theorem the relation $\leq_0$ of Example 2 is actually a *minimal* Borel-non-linearizable Borel order. (Compare with the “Glimm–Effros” theorem of Harrington, Kechris, Louveau [2] saying that E_0 is a minimal non-smooth Borel equivalence relation.)

¹ **Notation.** Several notions related to orders are sometimes understood differently, so let us take a space to fix an unambiguous meaning.

A binary relation $\preceq$ on a set X is a *partial quasi-order*, or *p. q.-o.* in brief, on X , iff $x \preceq y \wedge y \preceq z \implies x \preceq z$, and $x \preceq x$ for any $x \in X$. In this case, $\approx$ is the associated equivalence relation, *i. e.* $x \approx y$ iff $x \preceq y \wedge y \preceq x$.

If $x \approx x \implies x = x$ for any x then $\preceq$ is a *partial order*, or *p. o.* If in addition $x \preceq y \vee y \preceq x$ for all $x, y \in X$ then $\preceq$ is a *linear order* (l. o.).

Let $\preceq$ and $\preceq'$ be p. q.-o.'s on resp. X and X' . A map $h : X \longrightarrow X'$ will be called *half order preserving*, or *h. o. p.*, iff $x \preceq y \implies h(x) \preceq' h(y)$.

Finally a *linearization* is any h. o. p. map $h : \langle X; \preceq \rangle \longrightarrow \langle X'; \preceq' \rangle$, where $\preceq'$ is a l. o., satisfying $x \approx y \iff h(x) = h(y)$. $\bullet$

² If one enlarges $<_0$ so that, in addition, $a <_0 b$ whenever $a, b \in 2^\omega$ are such that $a(k) = 1$ and $b(k) = 0$ for all but finite k then the enlarged relation can be induced by a Borel action of $\mathbb{Z}$ on 2^ω , such that $a <_0 b$ iff $a = zb$ for some $z \in \mathbb{Z}$, $z > 0$.

Theorem 1 (Kanovei [11]) *Suppose that $\preceq$ is a Borel p. q.-o. on $\mathcal{N} = \omega^\omega$. Then exactly one of the following two conditions is satisfied:*

- (I^B) *$\preceq$ is Borel linearizable – moreover ³ in this case there are an ordinal $\alpha < \omega_1$ and a Borel linearization $h : \langle \mathcal{N}; \preceq \rangle \longrightarrow \langle 2^\alpha; \leq_{\text{lex}} \rangle$;*
- (II^B) *there exists a continuous h. o. p. 1 – 1 map $F : \langle 2^\omega; \leq_0 \rangle \longrightarrow \langle \mathcal{N}; \preceq \rangle$ such that $a \not\leq_0 b \implies F(a) \not\preceq F(b)$.*

Example 3. Let $\text{WO} = \{x \in \mathcal{N} : x \text{ codes an ordinal}\}$; for $x \in \text{WO}$ let $|x|$ be the ordinal coded by x . Define a Σ_1^1 p. q.-o. $x \preceq y$ iff either $y \notin \text{WO}$ or $x, y \in \text{WO}$ and $|x| \leq |y|$. (I^B) is impossible for $\preceq$ even via a non-Borel map h since orders $\langle 2^\alpha; \leq_{\text{lex}} \rangle$, $\alpha < \omega_1$, do not admit strictly increasing ω_1 -chains. (II^B) is also impossible via analytic maps F by the restriction theorem. Thus Theorem 1 fails for *analytic* relations. $\dashv$

Following ideas of Hjorth and Kechris [7], we involve longer orders, $2^{<\omega_1}$ and 2^{ω_1} , to match the nature of analytic p. q.-o.'s. A set $A \subseteq 2^{<\omega_1}$ will be called an *antichain* when it consists of pairwise $\subseteq$ -incomparable elements.

Theorem 2 *Suppose that $\preceq$ is a Σ_1^1 p. q.-o. on $\mathcal{N}$. Then at least one of the following two conditions is satisfied:*

- (I^A) *There is a linearization $h : \langle \mathcal{N}; \preceq \rangle \longrightarrow \langle 2^{\omega_1}; \leq_{\text{lex}} \rangle$ such that for any $\gamma < \omega_1$ the map $x \mapsto h(x) \upharpoonright \gamma$ is Borel, and has an ω_1 -Borel ⁴ code in $L[z]$ provided $\preceq$ is $\Sigma_1^1[z]$. In addition in each of the two ⁵ following cases there is an antichain $A \subseteq 2^{<\omega_1}$ and a Δ_2^1 in the codes linearization $h : \langle \mathcal{N}; \preceq \rangle \longrightarrow \langle A; \leq_{\text{lex}} \rangle$:*

- (a) *for any x the set $[x]_\approx = \{y : y \approx x\}$ is Borel;* ⁶
- (b) *the universe is a set generic ⁷ extension of a class $L[z_0]$, $z_0 \in \mathcal{N}$.*

(II^A) *As (II^B) of Theorem 1.*

³ Harrington *e. a.* [3] proved that any Borel l. o. is Borel order isomorphic to a l. o. $\langle X; \leq_{\text{lex}} \rangle$, where $X \subseteq 2^\alpha$ for some $\alpha < \omega_1$ and $\leq_{\text{lex}}$ is the lexicographical order.

⁴ Means: λ -Borel for an ordinal $\lambda < \omega_1$ which is not necessarily countable in $L[z]$.

⁵ An obvious parallel with the “Ulm classification” theorem in Hjorth and Kechris [7] allows to conjecture that the additional assertion is also true in the assumption of the existence of “sharps”, or an even weaker assumption in Friedman and Velickovic [1]. However the most interesting problem is to prove the additional assertion in **ZFC**.

⁶ This applies *e. g.* when $\preceq$ is a p. o.. Recall that $x \approx y$ iff $x \preceq y \wedge y \preceq x$.

⁷ Via any kind of set forcing. Compare with a theorem on thin Σ_1^1 equivalence relations in Hjorth [6].

Take notice that (I^A) and (II^A) here are compatible for instance in the assumption $V = L$. There possibly exist reasonable sufficient conditions (like: all Δ_2^1 sets are Lebesgue measurable) for (I^A) and (II^A) to be incompatible.

Applications for analytic equivalence relations

Theorem 2 applies for analytic equivalence relations viewed as a particular case of p. q.-o.'s.

Corollary 3 *Let E be a Σ_1^1 equivalence relation on $\mathcal{N}$. Then at least one of the following two conditions is satisfied:*

(I^E) *There is a map $h : \mathcal{N} \longrightarrow 2^{\omega_1}$ such that $x E y \iff h(x) = h(y)$ and for any $\gamma < \omega_1$ the map $x \mapsto h(x) \upharpoonright \gamma$ is Borel, and, provided $\preceq$ is $\Sigma_1^1[z]$, has an ω_1 -Borel code in $L[z]$. In addition in each of the two following cases there is an antichain $A \subseteq 2^{<\omega_1}$ and a Δ_2^1 in the codes map $h : \mathcal{N} \longrightarrow A$ such that $x E y \iff h(x) = h(y)$:*

- (a) *for any x the set $[x]_E = \{y : y E x\}$ is Borel;*
- (b) *the universe is a set generic extension of a class $L[z_0]$, $z_0 \in \mathcal{N}$.*

(II^E) *There exists a continuous 1 – 1 function $F : 2^\omega \longrightarrow \mathcal{N}$ such that $a E_0 b \iff F(a) E F(b)$.* □

This result, with (I^E) in the additional form, has been obtained by Hjorth and Kechris [7] in the subcase (I^E)(a) (as well as in the assumption of existence of sharps), by Friedman and Velickovic [1] in a hypothesis connected with weakly compact cardinals, and by Kanovei [10] in the subcase (I^E)(b).

(Recall that a map F as in (II^E) is called an *embedding* of E_0 in E – a continuous embedding in this case. A map h as in (I^E) is called a *reduction* of E to the equality on 2^{ω_1} or $2^{<\omega_1}$.)

Corollary 4 ⁸ *Assume that E is a $\Sigma_1^1[z]$ equivalence relation, $z \in \mathcal{N}$, and (II^E) of Corollary 3 fails. Then $x E y$ iff we have $x \in X \iff y \in X$ for every E -invariant Borel set $X \subseteq \mathcal{N}$ with an ω_1 -Borel code in $L[z]$.* □

Hjorth and Kechris [7] proved that any analytic E which does not satisfy (II^E) of Corollary 3 admits an effective reduction $h : \mathcal{N} \longrightarrow 2^{\omega_1}$ (i. e. we have $x E y \iff h(x) = h(y)$), however it is not clear whether the property mentioned in (I^E) of Corollary 3 holds for the reduction given in [7] and equally whether the reduction in [7] directly leads to Corollary 4.

⁸ Hjorth and Kechris told the author in April 1997 that they had known the result.

Organization of the proofs

The following theorem stands behind the results above.

Recall that if T is a tree on $\omega \times \omega \times \lambda$ then

$$[T] = \{ \langle x, y, f \rangle \in \mathcal{N}^2 \times \lambda^\omega : \forall m \ T(x \upharpoonright m, y \upharpoonright m, f \upharpoonright m) \}$$

and $\mathfrak{p}[T] = \{ \langle x, y \rangle : \exists f [T](x, y, f) \}$.

Theorem 5 *Let $\omega \leq \lambda < \omega_1$. Suppose that T and S are trees on $\omega \times \omega \times \lambda$ such that the sets $\preceq_T = \mathfrak{p}[T] \subseteq \preceq_S = \mathfrak{Cp}[S]$ are p. q.-o.'s on $\mathcal{N}$. Then at least one of the following two conditions is satisfied:*

- (I) *There are $\alpha < \omega_1$ and a ω_1 -Borel coded in $L[T, S]$ h. o. p. map $h : \langle \mathcal{N}; \preceq_T \rangle \longrightarrow \langle 2^\alpha; \leq_{\text{lex}} \rangle$ such that $h(x) = h(y) \implies x \approx_S y$.*
- (II) *There is a continuous $1 - 1$ h. o. p. map $F : \langle 2^\omega; \leq_0 \rangle \longrightarrow \langle \mathcal{N}; \preceq_T \rangle$ such that $a \not\leq_0 b \implies F(a) \not\leq_S F(b)$.*

The principal technical scheme of the proofs goes back to the papers of Harrington and Shelah [4], Shelah [13], and Horth [5] containing theorems on bi- κ -Souslin equivalence and order relations. However our version of the technique is free of any use of model theory including admissible sets.

On the other hand we exploit several technical achievements made in the study of the Borel orders (Harrington *e. a.* [3], Louveau [12]) by means of the Gandi – Harrington topology.

The two technical schemes, the one we use and the one based on the Gandi – Harrington topology, involve different kinds of “effective” sets in the forcing, but have many common points in the construction of the proofs (like a similar definition of the “regular” and “singular” cases, a similar construction of splitting systems *etc.*), although differ in many details.

As a matter of fact the Gandi – Harrington topology technique proves Theorem 1 shorter than we do here (see Kanovei [11]), but it has problems with the analytic case as it does not capture the proper type of effectiveness.

After some preliminaries in Section 1 (including an effective version of the classical separation theorem) we introduce the dichotomy in Section 2. Then the proof of Theorem 5 naturally develops itself in sections 3, 4, 5, and 6 (where we show that (II) of Theorem 5 is Shoenfield-absolute).

Theorem 2 (Section 7) will require a reflection argument saying that an analytic p. q.-o. has uncountably many indices for “upper” Borel approximations which are p. q.-o.'s, together with a delicate reasoning in the case of a generic universe, in Section 8.

1 Preliminaries

The proof of Theorem 5 is the major part of this paper.

We fix an ordinal λ , $\omega \leq \lambda < \omega_1$, and trees $T, S \subseteq (\omega \times \omega \times \lambda)^{<\omega}$.

Assume that both T and S are constructible.⁹

Suppose that $\preceq_T = \mathfrak{p}[T] \subseteq \preceq_S = \mathfrak{Cp}[S]$ are p. q.-o.'s on $\mathcal{N}$. Define $x \approx_S y$ iff $x \preceq_S y \wedge y \preceq_S x$ and $x \approx_T y$ similarly.

1.1 Coding Borel sets

We let $\mathcal{L}_{\lambda+1,0}$ be the infinitary language containing

- (i) constant symbols $\dot{x}, \dot{y}, \dot{z}, \dots$ for indefinite elements of $\mathcal{N} = \omega^\omega$ and constant symbols $\dot{f}, \dot{g}, \dots$ for indefinite elements of the set λ^ω ;
- (ii) elementary formulas of the form $\dot{x}(k) = l$ and $\dot{f}(k) = \alpha$, where $k, l \in \omega$ while $\alpha < \lambda$;
- (iii) conjunctions and disjunctions of size $\leq \lambda$, together with the ordinary propositional connectives, but it is assumed that any formula contains only finitely many constant symbols mentioned in (i).

(Quantifiers are not allowed). Thus formulas in $\mathcal{L}_{\lambda+1,0}$ code $(\lambda+1)$ -Borel subsets of spaces $\mathcal{N}^m \times (\lambda^\omega)^n$. For a formula, say, $\varphi(\dot{x}, \dot{f})$ we put

$$[\varphi] = \{\langle x, f \rangle \in \mathcal{N} \times \lambda^\omega : \varphi(x, f)\}.$$

For instance $[T](\dot{x}, \dot{y}, \dot{f})$ is a $\mathcal{L}_{\lambda+1,0}$ -formula; we shall denote it by $\dot{x} \preceq_{T,f} \dot{y}$. Similarly, the formula $\neg[S](\dot{x}, \dot{y}, \dot{f})$ will be denoted by $\dot{x} \preceq_{S,f} \dot{y}$. Formulas $\dot{x} \not\preceq_{S,f} \dot{y}$, $\dot{x} \approx_{S,f} \dot{y}$ etc. are derivatives. Then

$$x \preceq_T y \iff \exists f \in \lambda^\omega x \preceq_{T,f} y \quad \text{and} \quad x \preceq_S y \iff \forall f \in \lambda^\omega x \preceq_{S,f} y.$$

1.2 Consistency and separation

A formula φ is *consistent* if it has a model, *i. e.* becomes true after one suitably substitutes its constants by elements of $\mathcal{N}$ and λ^ω .

A *theory* in $\mathcal{L}_{\lambda+1,0}$ will be any set of formulas of $\mathcal{L}_{\lambda+1,0}$ containing a common (finite) list S of constants of type (i). (We shall usually consider *constructible* theories $\Phi \subseteq \mathcal{L}_{\lambda+1,0}$.) A theory is *consistent* if it has a model. A theory Φ is λ -*consistent* if every constructible subtheory $\Phi' \subseteq \Phi$ of cardinality $\leq \lambda$ in $\mathcal{L}$ is consistent.

⁹ Otherwise all entries of $\mathcal{L}$ from now on have to be uniformly changed to $\mathcal{L}[T, S]$.

A theory Φ λ -implies a formula ψ if $\Phi \cup \{\psi\}$ is λ -inconsistent. Other statements like this are to be understood accordingly.

The following theorem has a semblance of the Craig interpolation theorem, but essentially it belongs to the type of *separation* theorems.

Theorem 6 *Suppose that $\Phi(\dot{x}, \dot{y}, \dot{f}, \dots)$ and $\Psi(\dot{x}, \dot{y}', \dot{f}', \dots)$ are constructible theories in $\mathcal{L}_{\lambda+1,0}$ having $\dot{x}$ as the only common constant in the (finite) lists of constants. Assume that $\Phi(\dot{x}, \dot{y}, \dot{f}, \dots) \cup \Psi(\dot{x}, \dot{y}', \dot{f}', \dots)$ is λ -inconsistent. Then there is a $\mathcal{L}_{\lambda+1,0}$ -formula $\pi(\dot{x})$ λ -separating Φ from Ψ in the sense that $\Phi(\dot{x}, \dots)$ λ -implies $\pi(\dot{x})$ while $\Psi(\dot{x}, \dots)$ λ -implies $\neg \pi(\dot{x})$.*

Proof First of all we can assume that Φ and Ψ consist of single formulas, resp. $\varphi(\dot{x}, \dot{y}, \dot{f}, \dots)$ and $\psi(\dot{x}, \dot{y}', \dot{f}', \dots)$. Let, for the sake of simplicity, φ be $\varphi(\dot{x}, \dot{y})$ and ψ be $\psi(\dot{x}, \dot{f})$. Consider the sets

$$P = [\varphi] = \{\langle x, y \rangle : \varphi(x, y)\}, \quad Q = [\psi] = \{\langle x, f \rangle \in \mathcal{N} \times \lambda^\omega : \psi(x, f)\}.$$

The projections $X = \{x : \exists y P(x, y)\}$, $Y = \{x : \exists f Q(x, f)\}$ are disjoint Σ_1^1 sets by the inconsistency assumption, hence by the classical separation theorem they can be separated by a Borel set. Moreover as we demonstrated in Kanovei [9] (Theorem 7) in this case the separating set can be defined in the form $B = [\pi]$ for an appropriate $\mathcal{L}_{\lambda+1,0}$ -formula $\pi(\dot{x})$. $\square$

1.3 Hulls

By $\mathcal{F}(\dot{x})$ we shall denote the (constructible) collection of all $\mathcal{L}_{\lambda+1,0}$ -formulas $\varphi(\dot{x}) \in L$. For a theory $\Phi(\dot{x}, \dot{y}, \dots)$, $\mathcal{F}_{\dot{x}}[\Phi(\dot{x}, \dot{y}, \dots)]$ will be the set of all formulas $\varphi(\dot{x}) \in \mathcal{F}(\dot{x})$ which are λ -implied by $\Phi(\dot{x}, \dot{y}, \dots)$.

Lemma 7 *Suppose that $\Pi(\dot{x}, \dots)$ is a constructible theory in $\mathcal{L}_{\lambda+1,0}$ while $R(\dot{x}) \subseteq \mathcal{F}(\dot{x})$ is also constructible and satisfies $\mathcal{F}_{\dot{x}}[\Pi(\dot{x}, \dots)] \subseteq R(\dot{x})$. Then the theory $\Pi'(\dot{x}, \dots) = \Pi(\dot{x}, \dots) \cup R(\dot{x})$ satisfies $\mathcal{F}_{\dot{x}}[\Pi'(\dot{x}, \dots)] = \mathcal{F}_{\dot{x}}[R(\dot{x})]$.*

Proof Prove that $\mathcal{F}_{\dot{x}}[\Pi'(\dot{x}, \dots)] \subseteq \mathcal{F}_{\dot{x}}[R(\dot{x})]$ (the nontrivial direction). Let $\psi(\dot{x}) \in \mathcal{F}_{\dot{x}}[\Pi'(\dot{x}, \dots)]$. By definition there is a set $\Psi(\dot{x}) \in L$, $\Psi(\dot{x}) \subseteq R(\dot{x})$, of cardinality $\leq \lambda$ in L , such that $\Pi(\dot{x}, \dots) \cup \Psi(\dot{x})$ λ -implies $\psi(\dot{x})$. We conclude that the formula $(\bigwedge \Psi(\dot{x})) \implies \psi(\dot{x})$ belongs to $\mathcal{F}_{\dot{x}}[\Pi(\dot{x}, \dots)]$, hence to $R(\dot{x})$, which guarantees $\psi(\dot{x}) \in \mathcal{F}_{\dot{x}}[R(\dot{x})]$. $\square$

2 Monotone Borel functions and the dichotomy

To introduce the dichotomy we have to extend the language $\mathcal{L}_{\lambda+1,0}$ by Borel functions mapping $\mathcal{N}$ in a set of the form 2^α , where $\alpha < (\lambda^+)^L$. If such an α is fixed, let a *function code* be a sequence of the form $\vec{\varphi} = \langle \varphi_\gamma(\dot{x}) : \gamma < \alpha \rangle$ where each φ_γ is a $\mathcal{L}_{\lambda+1,0}$ -formula. Such a sequence defines a function $h_{\vec{\varphi}} : \mathcal{N} \longrightarrow 2^\alpha$ so that $h_{\vec{\varphi}}(x)(\gamma) = 1$ iff $\varphi_\gamma(x)$.

Define H_α to be the set of all h. o. p. maps $h_{\vec{\varphi}} : \langle \mathcal{N}; \preceq_T \rangle \longrightarrow \langle 2^\alpha; \leq_{\text{lex}} \rangle$, where $\vec{\varphi}$ is a *constructible* function code. Define $H = \bigcup_{\alpha < (\lambda^+)^L} H_\alpha$. Thus every function in H is an ω_1 -Borel (even $\lambda+1$ -Borel), coded in L , map from $\mathcal{N}$ to some 2^α , $\alpha < (\lambda^+)^L$, satisfying $x \preceq_T y \implies h(x) \leq_{\text{lex}} h(y)$.

But as a matter of fact functions in H will be used only via equalities of the form $h(x) = h(y)$ where $h = h_{\vec{\varphi}} \in H_\alpha$ for some $\alpha < (\lambda^+)^L$, viewed as shorthand for $\bigwedge_{\gamma < \alpha} (\varphi_\gamma(x) \iff \varphi_\gamma(y))$.

Let $\dot{x} \equiv_H \dot{y}$ be the theory $\{h(\dot{x}) = h(\dot{y}) : h \in H\}$. (Thus $\equiv_H$ defines an equivalence relation extending $\approx_T$.)

We have two cases.¹⁰

Case 1: the theory $\dot{x} \equiv_H \dot{y}$ λ -implies $\dot{x} \approx_{S,f} \dot{y}$.

Then clearly there is a single function $h \in H$ such that $h(\dot{x}) = h(\dot{y})$ already implies $\dot{x} \approx_{S,f} \dot{y}$. Then h satisfies (I) of Theorem 5.

Case 2: the theory $(\dot{x} \equiv_H \dot{y}) \cup \{\dot{x} \not\approx_{S,f} \dot{y}\}$ is λ -consistent.

Assuming this we shall work towards (II) of Theorem 5. We begin with a study of an important class of “conditionally downward closed” formulas.

Let $\mathcal{H}(\dot{x})$ be the (constructible as above) set of all formulas $\eta(\dot{x}) \in \mathcal{F}(\dot{x})$ satisfying the following: $\dot{x} \equiv_H \dot{x}'$ λ -implies $\eta(\dot{x}) \wedge \dot{x}' \preceq_{T,f} \dot{x} \implies \eta(\dot{x}')$.

Lemma 8 *Suppose that $\eta(\dot{x}) \in \mathcal{H}(\dot{x})$. Then there is a function $h \in H_{\alpha+1}$ for some $\alpha < (\lambda^+)^L$ such that $\eta(\dot{x}) \iff h(\dot{x}) = 0$. In particular the theory $\eta(\dot{x}) \wedge \neg \eta(\dot{y}) \wedge \dot{x} \equiv_H \dot{y}$ is then λ -inconsistent.*

Proof By definition there exists a function $h_0 \in H_\alpha$ for some $\alpha < (\lambda^+)^L$ satisfying $h_0(x) = h_0(y) \wedge \eta(x) \wedge x' \preceq_T x \implies \eta(x')$. Define $h(x) = h_0(x) \wedge 0$ whenever $\eta(x)$ and $h(x) = h_0(x) \wedge 1$ otherwise. $\square$

For a theory $\Phi(\dot{x}, \dot{y}, \dots)$, let $\mathcal{H}_{\dot{x}}[\Phi(\dot{x}, \dot{y}, \dots)] = \mathcal{F}_{\dot{x}}[\Phi(\dot{x}, \dot{y}, \dots)] \cap \mathcal{H}(\dot{x})$.

¹⁰ There is a point of dissatisfaction in the distribution on the two cases we use. It would be more natural to define Case 1 as that $\dot{x} \leq_H \dot{y}$ λ -implies $\dot{x} \preceq_S \dot{y}$, where $\dot{x} \leq_H \dot{y}$ is the theory $\{h(\dot{x}) \leq_{\text{lex}} h(\dot{y}) : h \in H\}$, which would improve (I) of Theorem 5 to the existence of a h. o. p. map satisfying $h(x) \leq h(y) \implies x \preceq_S y$. However then the arguments for Case 2, especially the key lemmas in the next section, do not go through.

3 The basic forcing

Let $\Xi(\dot{x})$ denote the (constructible) set of all formulas $\xi(\dot{x}) \in \mathcal{F}(\dot{x})$ which are λ -implied by the theory $\dot{x} \equiv_H \dot{y} \cup \{\dot{x} \not\prec_{S,f} \dot{y}\}$.

Let $\mathbf{P}$ be the set of all λ -consistent theories $\Pi(\dot{x}) \in \mathbf{L}$, $\Pi \subseteq \mathcal{F}(\dot{x})$, including $\Xi(\dot{x})$. Then $\mathbf{P} \in \mathbf{L}$, so we can view it as a forcing notion over $\mathbf{L}$.

Lemma 9 *Let $G \subseteq \mathbf{P}$ be $\mathbf{P}$ -generic over $\mathbf{L}$. Then there is a unique real $x = x_G \in \mathbf{L}[G]$ such that $\pi(\dot{x})$ holds in $\mathbf{L}[G]$ for any formula $\pi(\dot{x}) \in \bigcup G$.*

Proof Note that, for any n , the set D_n of all theories $\Pi(\dot{x}) \in \mathbf{P}$ which include $\dot{x}(n) = m$ for some m is dense in $\mathbf{P}$ and belongs to $\mathbf{L}$, hence $D_n \cap G \neq \emptyset$. The rest of the proof is standard. $\square$

3.1 Key lemmas

Lemma 10 *Let $\Pi(\dot{x})$ be a theory in $\mathbf{P}$. Then the theory $\Phi_\Pi(\dot{x}, \dot{y}, \dot{f}) =_{\text{df}} \Pi(\dot{x}) \cup \Pi(\dot{y}) \cup \dot{x} \equiv_H \dot{y} \cup \{\dot{x} \not\prec_{S,f} \dot{y}\}$ is λ -consistent and satisfies the equalities $\mathcal{F}_{\dot{x}}[\Phi_\Pi(\dot{x}, \dot{y}, \dot{f})] = \mathcal{F}_{\dot{x}}[\Pi(\dot{x})]$ and $\mathcal{F}_{\dot{y}}[\Phi_\Pi(\dot{x}, \dot{y}, \dot{f})] = \mathcal{F}_{\dot{y}}[\Pi(\dot{y})]$.*

Proof Let us first prove the consistency. Otherwise there is a formula $\pi(\dot{x}) \in \mathcal{F}_{\dot{x}}[\Pi(\dot{x})]$ and a function $h \in \mathbf{H}$ such that the formula $\pi(\dot{x}) \wedge \pi(\dot{y}) \wedge h(\dot{x}) = h(\dot{y}) \wedge \dot{x} \not\prec_{S,f} \dot{y}$ is λ -inconsistent.

The plan is to find functions $h', h'' \in \mathbf{H}$ such that the theories $\pi(\dot{x}) \wedge h'(\dot{x}) = h'(\dot{y}) \wedge \dot{y} \not\prec_{S,f} \dot{x}$ and $\pi(\dot{x}) \wedge h''(\dot{x}) = h''(\dot{y}) \wedge \dot{x} \not\prec_{S,f} \dot{y}$ are λ -inconsistent: then the formula $\neg \pi(\dot{x})$ belongs to Ξ , which is a contradiction because Π includes Ξ .

Consider the first theory (the other one is similar). By Lemma 8 it suffices to get a formula $\psi(\dot{x}) \in \mathcal{H}(\dot{x})$ such that $X = \lceil \pi \rceil \subseteq U = \lceil \psi \rceil$ and, for all $x \in X$ and $u \in U$, $h(x) = h(u)$ implies $u \preceq_S x$.

Let $Z = \{z : \forall x \in X (h(z) = h(x) \implies z \preceq_S x)\}$. Then $X \subseteq Z$ by the inconsistency assumption above.

Define a sequence of sets $X = X_0 \subseteq U_0 \subseteq X_1 \subseteq U_1 \subseteq \dots \subseteq Z$ and formulas $\varphi_n(\dot{x}) \in \mathcal{F}$ so that $U_n = \{u : \exists x \in X_n (h(x) = h(u) \wedge u \preceq_T x)\}$, $X_n = \lceil \varphi_n(\dot{x}) \rceil$, and the sequence of formulas φ_n is constructible.

Now, $\psi(\dot{x}) = \bigvee_n \varphi_n(\dot{x})$ is the formula required. (Note that $\lceil \psi \rceil = \bigcup_n X_n = \bigcup_n U_n$.) It remains to carry out the construction of X_n, U_n, φ_n .

Suppose that $X_n = \lceil \varphi_n(\dot{x}) \rceil \subseteq Z$ has been defined. Define U_n by the equality above. Then $X_n \subseteq U_n$, and $U_n \subseteq Z$. (Assume that $u \in U_n$, so

$u \preceq_T x$ for some $x \in X_n$ satisfying $h(x) = h(u)$. Take any $x' \in X$ such that $h(x') = h(u)$ and prove $u \preceq_S x'$. First of all $h(x) = h(x')$ hence $x \preceq_S x'$ because $x \in X_n \subseteq Z$. Now $u \preceq_S x'$ as $u \preceq_T x$.

Theorem 6 yields a formula $\beta(\dot{x}) \in \mathcal{F}$ such that the set $B = \lceil \beta \rceil$ satisfies $U_n \subseteq B \subseteq Z$. Take $X_{n+1} = B$ and $\varphi_{n+1} = \beta$.

Finally, as the choice of the formulas φ_n can be forced in L , the sequence of formulas can be chosen in L . This ends the proof of the consistency.

The equality $\mathcal{F}_{\dot{x}}[\Phi_{\Pi}(\dot{x}, \dot{y}, \dot{f})] = \mathcal{F}_{\dot{x}}[\Pi(\dot{x})]$ does not cause much trouble. Indeed suppose that $\pi(\dot{x}) \in \mathcal{F}(\dot{x}) \setminus \mathcal{F}_{\dot{x}}[\Pi(\dot{x})]$ (the nontrivial direction). Then the theory $\Pi'(\dot{x}) = \Pi(\dot{x}) \cup \{\neg \pi(\dot{x})\}$ is λ -consistent, hence belongs to $\mathbf{P}$. It follows from the above that $\Phi_{\Pi}(\dot{x}, \dot{y}, \dot{f}) \cup \{\neg \pi(\dot{x})\}$ is λ -consistent as well, hence $\pi(\dot{x}) \notin \mathcal{F}_{\dot{x}}[\Phi_{\Pi}(\dot{x}, \dot{y}, \dot{f})]$. $\square$

Lemma 11 *Assume that $\Pi(\dot{x})$ and $R(\dot{x})$ belong to $\mathbf{P}$, and $\mathcal{H}_{\dot{x}}[\Pi(\dot{x})] = \mathcal{H}_{\dot{x}}[R(\dot{x})]$. Then $\Psi_{\Pi R}(\dot{x}, \dot{y}, \dot{f}) =_{\text{df}} \Pi(\dot{x}) \cup R(\dot{y}) \cup \dot{x} \equiv_H \dot{y} \cup \{\dot{x} \preceq_{T,f} \dot{y}\}$ is a λ -consistent theory satisfying the equalities $\mathcal{F}_{\dot{x}}[\Psi_{\Pi R}(\dot{x}, \dot{y}, \dot{f})] = \mathcal{F}_{\dot{x}}[\Pi(\dot{x})]$ and $\mathcal{F}_{\dot{y}}[\Psi_{\Pi R}(\dot{x}, \dot{y}, \dot{f})] = \mathcal{F}_{\dot{y}}[R(\dot{y})]$.*

Proof As in the previous lemma, it suffices to prove the consistency. Suppose otherwise. Then there exist formulas $\pi(\dot{x}) \in \mathcal{F}_{\dot{x}}[\Pi(\dot{x})]$ and $\rho(\dot{y}) \in \mathcal{F}_{\dot{y}}[R(\dot{y})]$, and a function $h \in H$ such that the formula $\pi(\dot{x}) \wedge \rho(\dot{y}) \wedge h(\dot{x}) = h(\dot{y}) \wedge \dot{x} \preceq_{T,f} \dot{y}$ is inconsistent. In other words we have $x \not\preceq_T y$ whenever $x \in X = \lceil \pi \rceil$ and $y \in Y = \lceil \rho \rceil$ satisfy $h(x) = h(y)$.

Define $Z = \{z : \forall y \in Y (h(y) = h(z) \implies z \not\preceq_T y)\}$, so that $X \subseteq Z$ but $Y \cap Z = \emptyset$. The same iterated procedure as in the proof of Lemma 10 (with $U_n = \{u : \exists x \in X_n (h(x) = h(u) \wedge x \preceq_T u)\}$) yields a formula $\psi(\dot{x}) \in \mathcal{H}(\dot{x})$ such that the set $U = \lceil \psi \rceil$ satisfies $X \subseteq U \subseteq Z$. But this contradicts the assumption $\mathcal{H}_{\dot{x}}[\Pi(\dot{x})] = \mathcal{H}_{\dot{x}}[R(\dot{x})]$. $\square$

Corollary 12 *Suppose that $\Pi(\dot{x})$, $R(\dot{x})$ belong to $\mathbf{P}$ and $\mathcal{H}_{\dot{x}}[\Pi(\dot{x})] = \mathcal{H}_{\dot{x}}[R(\dot{x})]$. Then $U(\dot{x}, \dot{y}, \dot{f}) =_{\text{df}} \Pi(\dot{x}) \cup R(\dot{y}) \cup \dot{x} \equiv_H \dot{y} \cup \{\dot{x} \not\preceq_{S,f} \dot{y}\}$ is a λ -consistent theory satisfying the equalities $\mathcal{F}_{\dot{x}}[U(\dot{x}, \dot{y}, \dot{f})] = \mathcal{F}_{\dot{x}}[\Pi(\dot{x})]$ and $\mathcal{F}_{\dot{y}}[U(\dot{x}, \dot{y}, \dot{f})] = \mathcal{F}_{\dot{y}}[R(\dot{y})]$.*

Proof It suffices, as above, to prove the consistency. Suppose otherwise. Then the theory $\Phi_{\Pi}(\dot{x}, \dot{z}, \dot{f}) \cup \Psi_{R\Pi}(\dot{y}, \dot{z}, \dot{g})$ is λ -inconsistent as well. (Otherwise we have reals x, y, z satisfying $\Pi(x)$, $\Pi(z)$, $x \equiv_H z$, $R(y)$,

and $y \equiv_H z$, hence $x \equiv_H z$, and, in addition, $x \not\preceq_S z$ and $y \preceq_T z$, hence $x \not\preceq_S y$.) Theorem 6 yields a formula $\pi(\dot{z}) \in \mathcal{F}(\dot{z})$ λ -implied by $\Phi_\Pi(\dot{x}, \dot{z}, \dot{f})$ but λ -inconsistent with $\Psi_{R\Pi}(\dot{y}, \dot{z}, \dot{g})$, which is a contradiction as $\mathcal{F}_{\dot{z}}[\Phi_\Pi(\dot{x}, \dot{z}, \dot{f})] = \mathcal{F}_{\dot{z}}[\Pi(\dot{z})] = \mathcal{F}_{\dot{z}}[\Psi_{R\Pi}(\dot{y}, \dot{z}, \dot{g})]$ by lemmas 10, 11. $\square$

Corollary 13 *Suppose that $\Pi(\dot{x})$, $R(\dot{x})$ belong to $\mathbf{P}$ and $\mathcal{H}_{\dot{x}}[\Pi(\dot{x})] = \mathcal{H}_{\dot{x}}[R(\dot{x})]$. Then there are theories $\Pi'(\dot{x})$, $R'(\dot{x}) \in \mathbf{P}$ such that $\Pi \subseteq \Pi'$, $R \subseteq R'$, still $\mathcal{H}_{\dot{x}}[\Pi'(\dot{x})] = \mathcal{H}_{\dot{x}}[R'(\dot{x})]$, and $\Pi'(\dot{x}) \cup R'(\dot{x})$ is λ -inconsistent.*

Proof The theory $\Phi(\dot{x}, \dot{y}) =_{\text{df}} \Pi(\dot{x}) \cup R(\dot{x}) \cup (\dot{x} \equiv_H \dot{y}) \cup \{\dot{x} \neq \dot{y}\}$ is λ -consistent by the previous corollary. It easily follows that there exist numbers m and $k_x \neq k_y$ such that $\Phi'(\dot{x}, \dot{y}) =_{\text{df}} \Phi(\dot{x}, \dot{y}) \cup \{\dot{x}(m) = k_x\} \cup \{\dot{y}(m) = k_y\}$ is still λ -consistent. Now set $\Pi'(\dot{x}) = \mathcal{H}_{\dot{x}}[\Phi'(\dot{x}, \dot{y})]$ and $R'(\dot{y}) = \mathcal{H}_{\dot{y}}[\Phi'(\dot{x}, \dot{y})]$. $\square$

3.2 Two-dimensional modifications

There are two related forcing notions which produce pairs of reals.

Let $\mathbf{P}_{(2)}$ be the (constructible) set of all λ -consistent theories $\Delta(\dot{x}, \dot{y}) \in \mathbf{L}$ such that $\Xi(\dot{x}) \cup \Xi(\dot{y}) \subseteq \Delta(\dot{x}, \dot{y})$.

First modification. Recall that $\preceq_T = \mathbf{p}[T]$. The idea is to define a forcing which leads to pairs of reals satisfying $x \preceq_T y$.

We let ${}^T\mathbf{P}_{(2)}(\dot{x}, \dot{y}, \dot{f})$ be the set of all λ -consistent theories $\mathcal{T}(\dot{x}, \dot{y}, \dot{f})$ of the form $\Delta(\dot{x}, \dot{y}) \cup F \cup \dot{x} \equiv_H \dot{y} \cup \{\dot{x} \preceq_{T,f} \dot{y}\}$, where $\Delta \in \mathbf{P}_{(2)}$ and F is a finite collection of formulas $\dot{f}(k) = \alpha$ (where $k \in \omega$ and $\alpha < \lambda$).

For instance the theory $\Xi(\dot{x}) \cup \Xi(\dot{y}) \cup \dot{x} \equiv_H \dot{y} \cup \{\dot{x} \preceq_{T,f} \dot{y}\}$ (which is λ -consistent by Lemma 11) belongs to ${}^T\mathbf{P}_{(2)}$.

Lemma 14 *Let $G \subseteq {}^T\mathbf{P}_{(2)}$ be ${}^T\mathbf{P}_{(2)}$ -generic over $\mathbf{L}$. There is a unique triple $\langle x, y, f \rangle \in \mathbf{L}[G] \cap (\mathcal{N} \times \mathcal{N} \times \lambda^\omega)$ such that $\tau(x, y, f)$ holds for any formula $\tau(\dot{x}, \dot{y}, \dot{f}) \in \bigcup G$. In particular we have $x \preceq_T y$.*

Proof Analogous to Lemma 9. $\square$

Second modification. Recall that $\preceq_S = \mathbf{Cp}[S]$. Now the intension is to force pairs of reals $\langle x, y \rangle$ satisfying $x \not\preceq_S y$. We let ${}^S\mathbf{P}_{(2)}(\dot{x}, \dot{y}, \dot{f})$ be the set of all λ -consistent theories $\Sigma(\dot{x}, \dot{y}, \dot{f})$ of the form $\Delta(\dot{x}, \dot{y}) \cup F \cup \dot{x} \equiv_H \dot{y} \cup \{\dot{x} \not\preceq_{S,f} \dot{y}\}$, where F and Δ are as in the definition of ${}^T\mathbf{P}_{(2)}$.

To see that ${}^S\mathbf{P}_{(2)} \neq \emptyset$ note that $\Xi(\dot{x}) \cup \Xi(\dot{y}) \cup \dot{x} \equiv_H \dot{y} \cup \{\dot{x} \not\preceq_{S,f} \dot{y}\}$ is a λ -consistent theory by Lemma 10.

Lemma 15 *Let $G \subseteq {}^S\mathbf{P}_{(2)}$ be ${}^S\mathbf{P}_{(2)}$ -generic over $\mathbf{L}$. There is a unique triple $\langle x, y, f \rangle \in \mathbf{L}[G] \cap (\mathcal{N} \times \mathcal{N} \times \lambda^\omega)$ such that $\sigma(x, y, f)$ holds for any elementary formula $\sigma(\dot{x}, \dot{y}, \dot{f}) \in \bigcup G$. In particular we have $x \not\leq_S y$.*

Proof Analogous to Lemma 9. □

4 The product forcing

The forcing notion ${}^S\mathbf{P}_{(2)}$ executes too a tight control over generic reals. Fortunately generic $\leq_S$ -incomparable pairs can be obtained by another forcing, which connects the components in a much looser way, so that it is rather a kind of product forcing, with the factors equal to $\mathbf{P}$.

We let $\mathbf{P} \times_T \mathbf{P}$ be the set of all theories $\Upsilon(\dot{x}, \dot{z})$ of the form $\Pi(\dot{x}) \cup \mathbf{R}(\dot{z})$, where Π and $\mathbf{R}$ belong to $\mathbf{P}$ and satisfy $\mathcal{H}_{\dot{x}}[\Pi(\dot{x})] = \mathcal{H}_{\dot{x}}[\mathbf{R}(\dot{x})]$. The set $\mathbf{P} \times_T \mathbf{P}$ is constructible and non-empty.

Theorem 16 *Let $G \subseteq \mathbf{P} \times_T \mathbf{P}$ be $\mathbf{P} \times_T \mathbf{P}$ -generic over $\mathbf{V}$. There is a unique pair $\langle x, z \rangle \in \mathbf{V}[G] \cap \mathcal{N}^2$ such that $v(x, z)$ holds for any formula $v(\dot{x}, \dot{z}) \in \bigcup G$. Moreover we have $x \not\leq_S z$.*

Pairs $\langle x, z \rangle$ as in the theorem will be denoted by $\langle x_G, z_G \rangle$ and called $\mathbf{P} \times_T \mathbf{P}$ -generic over $\mathbf{V}$. ($\mathbf{V}$ is the universe of all sets as usual.)

Proof Let us concentrate on the proof that x_G and z_G are $\leq_S$ -incomparable; the rest is analogous to the above.

Suppose on the contrary that $x_G \leq_S z_G$ is $\mathbf{P} \times_T \mathbf{P}$ -forced over $\mathbf{V}$ by a “condition” $\Upsilon_0(\dot{x}, \dot{z}) = \Pi_0(\dot{x}) \cup \mathbf{R}_0(\dot{z}) \in \mathbf{P} \times_T \mathbf{P}$, where Π_0 and $\mathbf{R}_0$ belong to $\mathbf{P}_{(2)}$ and satisfy $\mathcal{H}_{\dot{x}}[\Pi_0(\dot{x})] = \mathcal{H}_{\dot{x}}[\mathbf{R}_0(\dot{x})]$.

We shall define a generic “rectangle” of reals x, z, x', z' , such that the following will be forced: $x \leq_S z$ and $x' \leq_S z'$ – by the forcing assumption, $z \leq_T x'$ – by Lemma 14, and $x \not\leq_S z'$ – by Lemma 15, which is a contradiction. The forcing $\mathfrak{P}$ used to get a required “rectangle” consists of forcing conditions of the following general form:

$$\mathfrak{p} = \langle \Upsilon(\dot{x}, \dot{z}), \mathcal{T}(\dot{z}, \dot{x}', \dot{f}), \Upsilon'(\dot{x}', \dot{z}'), \Sigma(\dot{x}, \dot{z}', \dot{f}) \rangle$$

such that the theories $\Upsilon(\dot{x}, \dot{z}) = \Pi(\dot{x}) \cup \mathbf{R}(\dot{z})$ and $\Upsilon'(\dot{x}', \dot{z}') = \Pi'(\dot{x}') \cup \mathbf{R}(\dot{z}')$ belong to $\mathbf{P} \times_T \mathbf{P}$, $\mathcal{T}$ belongs to ${}^T\mathbf{P}_{(2)}$, Σ belongs to ${}^S\mathbf{P}_{(2)}$, and

$$\begin{aligned} \Pi(\dot{x}) &= \mathcal{F}_{\dot{x}}[\Sigma(\dot{x}, \dot{z}', \dot{f})] & , & & \mathbf{R}(\dot{z}) &= \mathcal{F}_{\dot{z}}[\mathcal{T}(\dot{z}, \dot{x}', \dot{f})] \\ \Pi'(\dot{x}') &= \mathcal{F}_{\dot{x}'}[\mathcal{T}(\dot{z}, \dot{x}', \dot{f})] & , & & \mathbf{R}'(\dot{z}') &= \mathcal{F}_{\dot{z}'}[\Sigma(\dot{x}, \dot{z}', \dot{f})] \end{aligned}$$

Order $\mathfrak{P}$ componentwise: $\mathbf{p}_1$ is stronger than $\mathbf{p}_2$ iff $\Upsilon_2(\dot{x}, \dot{z}) \subseteq \Upsilon_1(\dot{x}, \dot{z})$, $\mathcal{T}_2(\dot{z}, \dot{x}', f) \subseteq \mathcal{T}_1(\dot{z}, \dot{x}', f)$, $\Upsilon'_2 \subseteq \Upsilon'_1$, and $\Sigma_2 \subseteq \Sigma_1$.

To get a condition in $\mathfrak{P}$ we start with the given theory $\Upsilon_0(x, z) = \Pi_0(\dot{x}) \cup R_0(\dot{z}) \in \mathbf{P} \times_T \mathbf{P}$. By definition $\mathcal{H}_{\dot{x}}[\Pi_0(\dot{x})] = \mathcal{H}_{\dot{x}}[R_0(\dot{x})]$.

It can be supposed that $\mathcal{F}_{\dot{x}}[\Pi_0(x)] = \Pi_0(x)$ (otherwise replace $\Pi_0(x)$ by $\mathcal{F}_{\dot{x}}[\Pi_0(\dot{x})]$) and $\mathcal{F}_{\dot{z}}[R_0(\dot{z})] = R_0(\dot{z})$.

The theory $\Sigma_0(\dot{x}, \dot{z}', f) =_{\text{df}} \Pi_0(\dot{x}) \cup R_0(\dot{z}') \cup \dot{x} \equiv_{\text{H}} \dot{z}' \cup \dot{x} \not\preceq_{S,f} \dot{z}'$ belongs to ${}^S\mathbf{P}_{(2)}$ and satisfies the equalities $\mathcal{F}_{\dot{x}}[\Sigma_0(\dot{x}, \dot{z}', f)] = \Pi_0(\dot{x})$ and $\mathcal{F}_{\dot{z}'}[\Sigma_0(\dot{x}, \dot{z}', f)] = R_0(\dot{z}')$ by Corollary 12. Similarly, by Lemma 11, the theory $\mathcal{T}_0(\dot{z}, \dot{x}', f) =_{\text{df}} R_0(\dot{z}) \cup \Pi_0(\dot{x}') \cup \dot{z} \equiv_{\text{H}} \dot{x}' \cup \dot{z} \preceq_{T,f} \dot{x}'$ belongs to ${}^T\mathbf{P}_{(2)}$ and satisfies $\mathcal{F}_{\dot{z}}[\mathcal{T}_0(\dot{z}, \dot{x}', f)] = R_0(\dot{z})$ and $\mathcal{F}_{\dot{x}'}[\mathcal{T}_0(\dot{z}, \dot{x}', f)] = \Pi_0(\dot{x}')$.

Now $\mathbf{p}_0 = \langle \Upsilon_0(\dot{x}, \dot{z}), \mathcal{T}_0(\dot{z}, \dot{x}', f), \Upsilon_0(\dot{x}', \dot{z}'), \Sigma_0(\dot{x}, \dot{z}', f) \rangle$ belongs to $\mathfrak{P}$.

Assertion Suppose that $\mathbf{p} = \langle \Upsilon, \mathcal{T}, \Upsilon', \Sigma \rangle \in \mathfrak{P}$, $\Upsilon_1(\dot{x}, \dot{z}) \in \mathbf{P} \times_T \mathbf{P}$, and $\Upsilon(\dot{x}, \dot{z}) \subseteq \Upsilon_1(\dot{x}, \dot{z})$. Then there exists $\mathbf{p}_1 = \langle \Upsilon_1, \mathcal{T}_1, \Upsilon'_1, \Sigma_1 \rangle \in \mathfrak{P}$ stronger than $\mathbf{p}$ (i. e. $\mathcal{T} \subseteq \mathcal{T}_1$, $\Upsilon' \subseteq \Upsilon'_1$, and $\Sigma \subseteq \Sigma_1$).

The same is true when we strengthen any of the other three components.

Proof By definition $\Upsilon_1(\dot{x}, \dot{z}) = \Pi_1(\dot{x}) \cup R_1(\dot{z})$ where Π_1 and R_1 belong to $\mathbf{P}_{(2)}$ and $\mathcal{H}_{\dot{x}}[\Pi_1(\dot{x})] = \mathcal{H}_{\dot{x}}[R_1(\dot{x})]$.

Let $\mathcal{T}_1(\dot{z}, \dot{x}', f) = \mathcal{T}(\dot{z}, \dot{x}', f) \cup \mathbf{R}_1(\dot{z})$. Lemma 7 yields $\mathcal{H}_{\dot{z}}[\mathcal{T}_1(\dot{z}, \dot{x}', f)] = \mathcal{H}_{\dot{z}}[\mathbf{R}_1(\dot{z})]$. Let $\Pi'_1(\dot{x}) = \mathcal{F}_{\dot{x}'}[\mathcal{T}_1(\dot{z}, \dot{x}', f)]$. Now $\mathcal{H}_{\dot{z}}[\Pi'_1(\dot{z})] = \mathcal{H}_{\dot{z}}[\mathbf{R}_1(\dot{z})]$ by Lemma 8 because even $\mathcal{T}(\dot{z}, \dot{x}', f)$ includes $\dot{z} \equiv_{\mathbf{H}} \dot{x}'$ by the definition of ${}^T\mathbf{P}_{(2)}$. Hence $\mathcal{H}_{\dot{x}}[\Pi'_1(\dot{x})] = \mathcal{H}_{\dot{x}}[\Pi_1(\dot{x})]$ by the above.

We define $R'_1(\dot{z}')$ using the other ark of the rectangle. Let $\Sigma_1(\dot{x}, \dot{z}', \dot{f}) = \Sigma(\dot{x}, \dot{z}', \dot{f}) \cup \Pi_1(\dot{x})$ and $R'_1(\dot{z}') = \mathcal{F}_{\dot{z}'}[\Sigma_1(\dot{x}, \dot{z}', \dot{f})]$. Then $\mathcal{H}_{\dot{x}}[R'_1(\dot{x})] = \mathcal{H}_{\dot{x}}[\Pi_1(\dot{x})]$ by lemmas 7 and 8 as above. Thus in particular we have $\mathcal{H}_{\dot{x}'}[R'_1(\dot{x}')] = \mathcal{H}_{\dot{x}'}[\Pi'_1(\dot{x}')]$, so that $\Upsilon'_1(\dot{x}', \dot{z}') = \Pi'_1(\dot{x}') \cup R'_1(\dot{z}')$ belongs to $\mathbf{P} \times_T \mathbf{P}$, closing the diagram. Clearly $\Upsilon \subseteq \Upsilon_1$. It easily follows from the construction that $\mathbf{p}_1 = \langle \Upsilon_1, \mathcal{T}_1, \Upsilon'_1, \Sigma_1 \rangle \in \mathfrak{P}$ is as required. $\dashv$

We continue the proof of Theorem 16. Consider a $\mathfrak{P}$ -generic extension $V[\mathfrak{G}]$ such that the generic set $\mathfrak{G} \subseteq \mathfrak{P}$ contains the condition $\mathfrak{p}_0$ defined above. It easily follows from the assertion just proved that $\mathfrak{G}$ results in a “rectangle” of reals $x, z, x', z' \in V[\mathfrak{G}]$ such that

- 1) The pair $\langle z, x' \rangle$ is ${}^T\mathbf{P}_{(2)}$ -generic over V , therefore we have $z \preccurlyeq_T x'$ in $V[\mathfrak{G}]$ by Lemma 14.

- 2) The pair $\langle x, z' \rangle$ is ${}^S\mathbf{P}_{(2)}$ -generic over V , therefore we have $x \not\preceq_S y$ in $V[\mathfrak{G}]$ by Lemma 15.
- 3) The pairs $\langle x, z \rangle$ and $\langle x', z' \rangle$ are $\mathbf{P} \times_T \mathbf{P}$ -generic over V , and moreover, the corresponding generic subsets of $\mathbf{P} \times_T \mathbf{P}$ contain the “condition” $\Upsilon_0(\dot{x}, \dot{z})$ fixed above; hence we have $x \preceq_S z$ and $x' \preceq_S z'$ in $V[\mathfrak{G}]$ by the choice of Υ_0 .

This is a contradiction because $\preceq_T \subseteq \preceq_S$ in $V[\mathfrak{G}]$ by absoluteness. $\square$

5 The construction of an embedding

We are going to define a continuous $1-1$ map $F : 2^\omega \longrightarrow \mathcal{N}$ satisfying (II) of Theorem 5. Our strategy will be to prove the existence of such a map in a κ -collapse generic extension V^+ of V , the universe of all sets, where κ is the cardinal $2^{2^{\aleph_0}}$ in V . This suffices to conclude that (II) of Theorem 5 holds in V by Lemma 18 of the next section.

5.1 Generic splitting family of theories

Let a *crucial pair* be any (ordered) pair $\langle u, v \rangle$ such that $u, v \in 2^m$ for some m and $u = 1^k \wedge 0^\wedge w$, $v = 0^k \wedge 1^\wedge w$ for some $k < m$ and $w \in 2^{m-k-1}$.

By the choice of V^+ the sets $\mathbf{P}$, ${}^T\mathbf{P}_{(2)}$, $\mathbf{P} \times_T \mathbf{P}$ have only V^+ -countably many subsets in V . Let $\{D(n) : n \in \omega\}$, $\{D^2(n) : n \in \omega\}$, $\{D_2(n) : n \in \omega\}$ be enumerations, in V^+ , of the collections of all dense (by *dense* we mean *open dense*) subsets of resp. $\mathbf{P}$, $\mathbf{P} \times_T \mathbf{P}$, ${}^T\mathbf{P}_{(2)}$. It will be assumed that each dense set has infinitely many indices in the relevant enumeration.

We shall define, in V^+ , a family of theories $\Pi_u(\dot{x}) \in \mathbf{P}$ (where $u \in 2^{<\omega}$) and $\mathcal{T}_{uv}(\dot{x}, \dot{y}, \dot{f}) \in {}^T\mathbf{P}_{(2)}$ (where $\langle u, v \rangle$ is a crucial pair in some 2^n) satisfying the following conditions, for all $u \in 2^{<\omega}$ and $i = 0, 1$:

- (i) $\Pi_u \in D(n)$ whenever $u \in 2^n$; $\Pi_u(\dot{x}) \subseteq \Pi_{u \wedge i}(\dot{x})$;
- (ii) if $\langle u, v \rangle$ is a crucial pair in 2^n then $\mathcal{T}_{uv} \in D_2(n)$; $\mathcal{T}_{uv}(\dot{x}, \dot{y}, \dot{f}) \subseteq \mathcal{T}_{u \wedge i, v \wedge i}(\dot{x}, \dot{y}, \dot{f})$;
- (iii) if $u, v \in 2^n$ and $u(n-1) \neq v(n-1)$ then $\Pi_u(\dot{x}) \cup \Pi_v(\dot{z}) \in \mathbf{P} \times_T \mathbf{P}$, moreover $\in D^2(n)$, and $\Pi_u(\dot{x}) \cup \Pi_v(\dot{x})$ is λ -inconsistent;
- (iv) $\mathcal{F}_{\dot{x}}[\mathcal{T}_{uv}(\dot{x}, \dot{y}, \dot{f})] = \Pi_u(\dot{x})$ and $\mathcal{F}_{\dot{y}}[\mathcal{T}_{uv}(\dot{x}, \dot{y}, \dot{f})] = \Pi_v(\dot{y})$ — then in particular $\mathcal{F}_{\dot{x}}[\Pi_u(\dot{x})] = \Pi_u(\dot{x})$ for all u .

Remark 17 Since theories in ${}^T\mathbf{P}_{(2)}$ contain $\dot{x} \equiv_{\mathbf{H}} \dot{y}$, it follows from (iv) by Lemma 8 that $\mathcal{H}_{\dot{x}}[\Pi_u(\dot{x})] = \mathcal{H}_{\dot{x}}[\Pi_v(\dot{x})]$ for all crucial pairs $\langle u, v \rangle$. Therefore $\mathcal{H}_{\dot{x}}[\Pi_u(\dot{x})] = \mathcal{H}_{\dot{x}}[\Pi_v(\dot{x})]$ for all $u, v \in 2^n$ and $n \in \omega$ as any two tuples $u, v \in 2^n$ are connected by a (unique) chain of crucial pairs. $\square$

Let us first of all demonstrate that the existence of such a system yields a continuous map F in V^+ which witnesses (II) of Theorem 5.

Lemma 9 and (i) imply that for any $a \in 2^\omega$ there is a unique real, denoted by $F(a)$, satisfying every formula in $\bigcup_{n \in \omega} \Pi_{a \upharpoonright n}(\dot{x})$, and the map F is continuous. Moreover F is 1 – 1 by (iii).

Suppose that $a, b \in 2^\omega$ and $a \not\leq_0 b$, so that $a(n) \neq b(n)$ for infinitely many n . It follows then from (iii) and Theorem 16 that $F(a) \not\leq_S F(b)$.

Let us check that F satisfies (II) of Theorem 5. Suppose that $a, b \in 2^\omega$ are $\leq_0$ -neighbours, *i. e.* $a = 1^k \wedge 0^\omega$ and $b = 0^k \wedge 1^\omega$ for some $k \in \omega$ and $c \in 2^\omega$. Then $\langle a \upharpoonright n, b \upharpoonright n \rangle$ is a crucial pair for all $n > k$. Therefore, by (ii) and Lemma 14, there is a unique triple $\langle x, y, f \rangle \in \mathcal{N}^2 \times \lambda^\omega$ which satisfies every formula in $\bigcup_{n \in \omega} \mathcal{T}_{a \upharpoonright n, b \upharpoonright n}(\dot{x}, \dot{y}, \dot{f})$, and now $x = F(a)$, $y = F(b)$ by (iv). This implies $F(a) \leq_T F(b)$ by (ii).

5.2 The construction of theories

We argue in V^+ .

To define Π_Λ (where Λ is the empty sequence, the only member of 2^0) consider first the theory $\Xi(\dot{x})$, see Subsection 3. As clearly $\Xi(\dot{x}) \in \mathbf{P}$, there is a theory $\Pi(\dot{x}) \in D(0)$ including $\Xi(\dot{x})$. Let $\Pi_\Lambda(\dot{x}) = \Pi(\dot{x})$.

Suppose that the construction has been completed up to a level n , and expand it to the next level.

To start with we set $\Pi_{s \wedge i}(\dot{x}) = \Pi_s(\dot{x})$ for all $s \in 2^n$ and $i = 0, 1$, and $\mathcal{T}_{s \wedge i, t \wedge i}(\dot{x}, \dot{y}, \dot{f}) = \mathcal{T}_{st}(\dot{x}, \dot{y}, \dot{f})$ whenever $i = 0, 1$ and $\langle s, t \rangle$ is a crucial pair in 2^n . For the “initial” pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$, let $\mathcal{T}_{1^n \wedge 0, 0^n \wedge 1}$ be the theory $\Pi_{0^n}(\dot{x}) \cup \Pi_{0^n}(\dot{y}) \cup (\dot{x} \equiv_{\mathbf{H}} \dot{y}) \cup \{\dot{x} \leq_{T, f} \dot{y}\}$. Then, by Lemma 11, $\mathcal{T}_{1^n \wedge 0, 0^n \wedge 1} \in {}^T\mathbf{P}_{(2)}$ and $\mathcal{F}_{\dot{x}}[\mathcal{T}_{1^n \wedge 0, 0^n \wedge 1}(\dot{x}, \dot{y}, \dot{f})] = \Pi_{1^n \wedge 0}(\dot{x})$, $\mathcal{F}_{\dot{y}}[\mathcal{T}_{1^n \wedge 0, 0^n \wedge 1}(\dot{x}, \dot{y}, \dot{f})] = \Pi_{0^n \wedge 1}(\dot{y})$.

This ends the definition of “initial values” at the $n+1$ -th level. The plan is to gradually enforce the theories in order to fulfill the requirements.

Step 1. Take care of item (i). Consider an arbitrary $u_0 = s_0 \wedge i \in 2^{n+1}$. As $D(n)$ is dense there is a theory $\Pi'(\dot{x}) \in D(n)$ including $\Pi_{u_0}(\dot{x})$. We can assume that $\Pi'(\dot{x}) = \mathcal{F}_{\dot{x}}[\Pi'(\dot{x})]$ for otherwise change $\Pi'(\dot{x})$ by $\mathcal{F}_{\dot{x}}[\Pi'(\dot{x})]$.

The intension is to take $\Pi'(\dot{x})$ as the “new” Π_{u_0} . But this change has to be expanded through the net of crucial pairs, in order to preserve (iv). (Fortunately the tree of all crucial pairs in 2^{n+1} is a chain.)

Thus put $\Pi'_{u_0}(\dot{x}) = \Pi'(\dot{x})$. Suppose that Π'_u has been defined, includes Π_u , the older version, and satisfies $\mathcal{F}_{\dot{x}}[\Pi'_u(\dot{x})] = \Pi'_u(\dot{x})$, for some $u \in 2^{n+1}$ which is connected by a crucial pair with a not yet encountered $v \in 2^{n+1}$. Define $\mathcal{T}'_{uv}(\dot{x}, \dot{y}, \dot{f})$ to be $\Pi'_u(\dot{x}) \cup \mathcal{T}_{uv}(\dot{x}, \dot{y}, \dot{f})$ and $\Pi'_v(\dot{y})$ to be $\mathcal{F}_{\dot{y}}[\mathcal{T}'_{uv}(\dot{x}, \dot{y}, \dot{f})]$. Note that $\Pi'_v(\dot{y})$ includes $\Pi_v(\dot{y})$ because (iv) is assumed for the old theories $\Pi_u, \Pi_v, \mathcal{T}_{uv}$. Note that (iv) holds for the new theories $\Pi'_u, \Pi'_v, \mathcal{T}'_{uv}$: indeed $\mathcal{F}_{\dot{x}}[\mathcal{T}'_{uv}(\dot{x}, \dot{y}, \dot{f})] = \Pi'_u(\dot{x})$ follows from Lemma 7.

The construction describes how the change from Π_{u_0} to Π'_{u_0} spreads through the chain of crucial pairs in 2^{n+1} , resulting in a system of new theories, Π'_u and $\mathcal{T}'_{uv}$, which satisfy (i) for the particular $u_0 \in 2^{n+1}$.

We iterate this construction consecutively for all $u_0 \in 2^{n+1}$, getting finally a system of theories satisfying (i) (fully) (and (iv)) which we shall denote by Π_u and $\mathcal{T}_{uv}$ from now on.

Step 2. Take care of item (iii). Let us fix a pair of u_0 and v_0 in 2^{n+1} , such that $u_0(n) = 0$ and $v_0(n) = 1$. By the density of $D^2(n)$, there is a theory $\Pi'_{u_0}(\dot{x}) \cup \Pi'_{v_0}(\dot{y}) \in D^2(n)$ which includes $\Pi_{u_0}(\dot{x}) \cup \Pi_{v_0}(\dot{y})$. We may assume that $\Pi'_{u_0}(\dot{x}) = \mathcal{F}_{\dot{x}}[\Pi'_{u_0}(\dot{x})]$ and $\Pi'_{v_0}(\dot{y}) = \mathcal{F}_{\dot{y}}[\Pi'_{v_0}(\dot{y})]$. We can also assume, by Corollary 13, that $\Pi'_{u_0}(\dot{x}) \cup \Pi'_{v_0}(\dot{x})$ is λ -inconsistent.

Spread the change from Π_{u_0} to Π'_{u_0} and from Π_{v_0} to Π'_{v_0} through the chain of crucial pairs in 2^{n+1} until the two waves of spreading meet each other at the pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$. This leads to a system of theories Π'_u and $\mathcal{T}'_{uv}$ which satisfy (iii) for the particular pair $\langle u_0, v_0 \rangle$ and still satisfy (iv) with the exception of the “meeting” crucial pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$ (for which basically $\mathcal{T}'_{1^n \wedge 0, 0^n \wedge 1}$ is not yet defined for this step).

Take notice that the construction of Step 1 has left $\mathcal{T}_{1^n \wedge 0, 0^n \wedge 1}$ in the form $\Pi_{1^n \wedge 0}(\dot{x}) \cup \Pi_{0^n \wedge 1}(\dot{y}) \cup (\dot{x} \equiv_{\mathbf{H}} \dot{y}) \cup \{\dot{x} \preceq_{T, f} \dot{y}\}$ (where $\Pi_{1^n \wedge 0}$ and $\Pi_{0^n \wedge 1}$ are the “versions” at the end of Step 1). We now have new λ -consistent theories, $\Pi'_{1^n \wedge 0}$ and $\Pi'_{0^n \wedge 1}$, including resp. $\Pi_{1^n \wedge 0}$ and $\Pi_{0^n \wedge 1}$ and satisfying $\mathcal{H}_{\dot{x}}[\Pi'_{0^n \wedge 1}(\dot{x})] = \mathcal{H}_{\dot{x}}[\Pi'_{0^n \wedge 1}(\dot{x})]$. (See Remark 17; recall that $\mathcal{H}_{\dot{x}}[\Pi'_{u_0}] = \mathcal{H}_{\dot{x}}[\Pi'_{v_0}]$ for the initial pair simply because $\Pi'_{u_0}(\dot{x}) \cup \Pi'_{v_0}(\dot{y}) \in \mathbf{P} \times_T \mathbf{P}$.) Now the theory $\Pi'_{1^n \wedge 0}(\dot{x}) \cup \Pi'_{0^n \wedge 1}(\dot{y}) \cup (\dot{x} \equiv_{\mathbf{H}} \dot{y}) \cup \{\dot{x} \preceq_{T, f} \dot{y}\}$ taken as $\mathcal{T}'_{1^n \wedge 0, 0^n \wedge 1}$ belongs to ${}^T\mathbf{P}_{(2)}$ and satisfies (iv) for the pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$ by Lemma 11. This ends the consideration of the pair $\langle u_0, v_0 \rangle$.

Applying this construction consecutively for all pairs of $u_0 \in P_0$ and $v_0 \in P_1$ (including the pair $\langle 1^n \wedge 0, 0^n \wedge 1 \rangle$) we finally get a system of theories satisfying (i), (iii), and (iv), which will be denoted still by Π_u and $\mathcal{T}_{uv}$.

Step 3. We finally take care of (ii). Consider a particular crucial pair $\langle u_0, v_0 \rangle$ in 2^{n+1} . By the density of $D_2(n)$, there is a theory $\mathcal{T}'_{u_0, v_0}(\dot{x}, \dot{y}, \dot{f})$ in $D_2(n)$ including $\mathcal{T}_{u_0, v_0}(\dot{x}, \dot{y}, \dot{f})$.

Define $\Pi'_{u_0}(\dot{x}) = \mathcal{F}_{\dot{x}}[\mathcal{T}'_{u_0, v_0}(\dot{x}, \dot{y}, \dot{f})]$ and $\Pi'_{v_0}(\dot{y}) = \mathcal{F}_{\dot{y}}[\mathcal{T}'_{u_0, v_0}(\dot{x}, \dot{y}, \dot{f})]$ and spread this change through the chain of crucial pairs in 2^{n+1} . (Note that $\mathcal{H}_{\dot{x}}[\Pi'_{u_0}(\dot{x})] = \mathcal{H}_{\dot{x}}[\Pi'_{v_0}(\dot{x})]$ because theories in ${}^T\mathbf{P}_{(2)}$ include $\dot{x} \equiv_{\mathbf{H}} \dot{y}$. This implies $\mathcal{H}_{\dot{x}}[\Pi'_u(\dot{x})] = \mathcal{H}_{\dot{x}}[\Pi'_v(\dot{x})]$ for all $u, v \in 2^{n+1}$ after the spreading.)

Executing this construction for all crucial pairs in 2^{n+1} we finally end the construction, in V^+ , of a system of theories satisfying (i) through (iv).

□ (Theorem 5)

6 Why embedding $\leq_0$ is absolute

The aim of this section is to prove that (II) of Theorem 5 and (II^A) of Theorem 2 are absolute statements. By the way this fills the gap left in the proof of Theorem 5 (see the beginning of Section 5).

Lemma 18 *If $p \in \mathcal{N}$ and $\preceq$ is a $\Sigma_1^1(p)$ p. q.-o. then (II^A) of Theorem 2 is equivalent to a $\Sigma_2^1(p)$ statement uniformly in p . Similarly if $\preceq_T \subseteq \preceq_S$ are resp. $\Sigma_1^1(p)$ p. q.-o. and $\Pi_1^1(p)$ p. q.-o. then (II) of Theorem 5 is equivalent to a $\Sigma_2^1(p)$ statement uniformly in p .*

Proof ¹¹ We consider only (II) of Theorem 5; the other statement is pretty similar. The aim does not seem easy: at the first look the statement is Σ_3^1 . To reduce it to Σ_2^1 we use *Borel approximations* of $\preceq_T$ and $\preceq_S$.

Recall that $\mathbf{WO} = \{z \in \mathcal{N} : z \text{ codes an ordinal}\}$; for $z \in \mathbf{WO}$ let $|z|$ be the ordinal coded by z , and $\mathbf{WO}_\nu = \{z \in \mathbf{WO} : |z| = \nu\}$.

Being a Σ_1^1 subset of $\mathcal{N}^2$, the relation $\preceq_T$ classically has the form $\preceq_T = \bigcup_{\nu < \omega_1} \preceq_{T, \nu}$ where $\langle \preceq_{T, \nu} : \nu < \omega_1 \rangle$ is an increasing sequence of Borel subsets of $\preceq_T$. Moreover there is a Π_1^1 formula $\pi(z, x, y)$ (containing p as a parameter) such that $x \preceq_{T, \nu} y \iff \pi(z, x, y)$ whenever $z \in \mathbf{WO}_\nu$. (There also exists a Σ_1^1 formula with the same property.)

Similarly $\preceq_S = \bigcap_{\nu < \omega_1} \preceq_{S, \nu}$, where $\langle \preceq_{S, \nu} : \nu < \omega_1 \rangle$ is an increasing sequence of Borel supersets of $\preceq_S$, and there is a Σ_1^1 formula $\sigma(z, x, y)$ such that $x \preceq_{S, \nu} y \iff \sigma(z, x, y)$ whenever $z \in \mathbf{WO}_\nu$.

The following statement is clearly $\Sigma_2^1(p)$ (use formulas π and σ):

- (II') *There is a continuous 1 – 1 map $F' : 2^\omega \longrightarrow \mathcal{N}$ and a countable ordinal ν such that*
- (a) $a \leq_0 b$ *implies* $F'(a) \preceq_{T, \nu} F'(b)$;
 - (b) $a \not\leq_0 b$ *implies* $\neg F'(a) \preceq_{S, \nu} F'(b)$.

¹¹ The proof involves an idea communicated to the author by G. Hjorth with a reference to Hjorth and Kechris [7], Section 3, where the idea is realized in terms of category.

Thus it remains to prove that (II) of Theorem 5 is equivalent to (II'). The hard part is to prove that (II) implies (II'). To prove this direction we consider a κ -collapse generic extension V^+ of V , the universe of all sets, where κ is $2^{\aleph_0}$ in V . As (II) is $\Sigma_3^1(p)$ while (II') is $\Sigma_2^1(p)$, it suffices to prove that (II) implies (II') in V^+ .

We can enumerate in V^+ by natural numbers all dense subsets of $2^{<\omega}$ and $2^{<\omega} \times 2^{<\omega}$ (the Cohen forcing and its square) which belong to V . This allows to define in V^+ infinite sequences $\langle u_n : n \in \omega \rangle$ and $\langle v_n : n \in \omega \rangle$ such that $u_n, v_n \in 2^{l(n)}$ for some $l(n)$ for all n , and for any n :

- (*) if $u, v \in 2^l$ where $l = n + \sum_{m=0}^{n-1} l(m)$ then the pairs $\langle u \wedge u_n, v \wedge v_n \rangle$ and $\langle u \wedge v_n, v \wedge u_n \rangle$ belong to the n -th dense subset of $2^{<\omega} \times 2^{<\omega}$.

Define in V^+ , for each $a \in 2^\omega$, $G(a) = w_0 \wedge w_1 \wedge w_2 \wedge \dots$, where $w_n = u_n$ whenever $a(n) = 0$ and $w_n = v_n$ whenever $a(n) = 1$. Then G is continuous and $1-1$, therefore the map $F'(a) = F(G(a))$ is continuous and $1-1$ as well. (Here F is a map which witnesses (II) of Theorem 5 in V^+ .) Prove that F' witnesses (II').

Suppose that $a, b \in 2^\omega$ and $a \leq_0 b$. Then by definition both $a' = G(a)$ and $b' = G(b)$ are **Cohen** generic over V and $a' \leq_0 b'$, $F(a') \preceq_T F(b')$ (by the choice of F), even in $V[a', b']$, which implies $F(a') \preceq_{T, \nu} F(b')$ for an appropriate $\nu < \omega_1^{V[a', b']}$. Since the difference between a' and b' is finite the latter statement is a property of a' , hence it is **Cohen** forced over V . It follows that there is a countable in $V[a']$ (hence in V^+) ordinal ν_T such that we have, in V^+ , $F'(a) \preceq_{T, \nu_T} F'(b)$ whenever $a, b \in 2^\omega$ satisfy $a \leq_0 b$.

Suppose that $a, b \in 2^\omega$ and $a \not\leq_0 b$. Then by definition $\langle G(a), G(b) \rangle$ is **Cohen**² generic over V , in particular $a' = G(a)$ and $b' = G(b)$ satisfy $a' \not\leq_0 b'$, therefore $F(a') \not\preceq_S F(b')$, which implies $\neg F(a') \preceq_{S, \nu} F(b')$ for an ordinal $\nu < \omega_1^{V[a', b']}$. As above there is a single ordinal $\nu_S < \omega_1^{V^+}$ such that we have, in V^+ , $\neg F'(a) \preceq_{S, \nu_S} F'(b)$ whenever $a \not\leq_0 b$.

It remains to take $\nu = \max \{ \nu_T, \nu_S \}$. □

7 Borel and analytic order relations

This section proves theorems 1 and 2, with the exception of the additional statement in (I^A) of Theorem 2.

7.1 Borel orders: Theorem 1

Let $\preceq$ be a Borel p. q.-o. on $\mathcal{N}$. In view of Theorem 5 it suffices to prove that (I^B) and (II^B) of Theorem 1 are incompatible. *Suppose otherwise.*

The superposition of the maps F and h is then a Borel h. o. p. map $\phi : \langle 2^\omega; \leq_0 \rangle \longrightarrow \langle 2^\alpha; \leq_{\text{lex}} \rangle$ satisfying the following: $\phi(a) = \phi(b)$ implies that $a \mathbf{E}_0 b$, i. e. a and b are $\leq_0$ -comparable.

Therefore, as any $\mathbf{E}_0$ -class is $\leq_0$ -ordered as $\mathbb{Z}$, ω , or ω^* , the ϕ -image $X_a = \phi'' [a]_{\mathbf{E}_0}$ of the $\mathbf{E}_0$ -class of any $a \in 2^\omega$ is $\leq_{\text{lex}}$ -ordered as a subset of $\mathbb{Z}$. If $X_a = \{x_a\}$ is a singleton then put $\psi(a) = x_a$.

Assume that X_a contains at least two points. In this case we can effectively pick an element in X_a ! Indeed there is a maximal sequence $u \in 2^{<\alpha}$ such that $u \subset x$ for each $x \in X_a$. Then the set $X_a^{\text{left}} = \{x \in X : u^\frown 0 \subset x\}$ contains the $\leq_{\text{lex}}$ -largest element, which we denote by $\psi(a)$.

To conclude ψ is a Borel reduction of $\mathbf{E}_0$ to the equality on 2^α , i. e. $a \mathbf{E}_0 b$ iff $\psi(a) = \psi(b)$, which is impossible.

□ (*Theorem 1*)

7.2 The general case of analytic relations

Consider an analytic p. q.-o. $\preceq$ on $\mathcal{N}$. We shall w. l. o. g. assume that $\preceq$ is Σ_1^1 , so that $\preceq = \preceq_T = \mathbf{p}[T]$, where T is a recursive tree in $(\omega \times \omega \times \omega)^{<\omega}$. We also suppose that

(†) $\preceq_T$ does not satisfy (II^A) of Theorem 2.

The aim is to prove that then $\preceq_T$ satisfies (I^A) (still leaving apart the additional part) of Theorem 2.

Recall that H_α is defined in Section 2. Let $H^* = \bigcup_{\alpha < \omega_1} H_\alpha$; this includes the set H also defined in Section 2. By definition each $h \in H^*$ is a Borel function of certain type, coded in L . Let us fix a *constructible in the codes* enumeration $H^* = \{h_\alpha : \alpha < \omega_1\}$. Define the concatenation

$$\mathbf{h}(x) = h_0(x)^\frown h_1(x)^\frown \dots^\frown h_\alpha(x)^\frown \dots \quad (\alpha < \omega_1).$$

Prove that $\mathbf{h} : \langle \mathcal{N}; \preceq_T \rangle \longrightarrow \langle 2^{<\omega_1}; \leq_{\text{lex}} \rangle$ is a linearization. First of all $\mathbf{h}$ is a h. o. p. map from $\langle \mathcal{N}; \preceq_T \rangle$ to $\langle 2^{<\omega_1}; \leq_{\text{lex}} \rangle$ because each h_α is h. o. p. by definition. Thus it remains to prove that $\mathbf{h}(x) = \mathbf{h}(y) \implies x \approx_T y$.

This involves a *reflection lemma* for analytic p. q.-o.'s.

Being a Σ_1^1 set, $\preceq_T$ has the form $\preceq_T = \bigcap_{\nu < \omega_1} \preceq_T^\nu$, where each $\preceq_T^\nu$ is an ω_1 -Borel set coded in L and $\preceq_T^\mu \subseteq \preceq_T^\nu$ provided $\nu < \mu$. In addition we have the following **boundedness principle**: if $\preceq_T \subseteq X$, where $X \subseteq \mathcal{N}^2$ is a Π_1^1 set, then there is $\nu < \omega_1$ such that $\preceq_T^\nu \subseteq X$.

Take notice that the sets $\preceq_T^\nu$ are not necessarily p. q.-o.'s. However

Lemma 19 Assume that $B \subseteq \mathcal{N}^2$ is a Borel set and $\preceq_T \subseteq B$. Then there is $\nu < \omega_1$ such that $\preceq_T^\nu \subseteq B$ and $\preceq_T^\nu$ is a p. q.-o..¹²

Proof Let us first prove a weaker statement: there is $\mu < \omega_1$ such that

$$x \preceq_T^\mu y \implies \forall x' \forall y' (x' \preceq_T x \wedge y \preceq_T y' \implies x' \preceq_T^\mu y').$$

Note that by the boundedness there exists an ordinal $\mu_0 < \omega_1$ such that $\preceq_T^{\mu_0} \subseteq B$. Suppose that an ordinal $\mu_n \geq \mu_0$ has been defined. Put $C(x, y)$ iff $\forall x' \forall y' (x' \preceq_T x \wedge y \preceq_T y' \implies x' \preceq_T^{\mu_n} y')$, so that C is a Π_1^1 set and $\preceq_T \subseteq C \subseteq \preceq_T^{\mu_n}$. Using the boundedness principle again we get an ordinal $\mu_{n+1} \geq \mu_n$ satisfying $\preceq_T \subseteq \preceq_T^{\mu_{n+1}} \subseteq C$, so that by definition

$$x \preceq_T^{\mu_{n+1}} y \implies \forall x' \forall y' (x' \preceq_T x \wedge y \preceq_T y' \implies x' \preceq_T^{\mu_n} y').$$

It remains to define $\mu = \sup_n \mu_n$.

Starting the proof of the lemma, we choose ν_0 so that $\preceq_T^{\nu_0} \subseteq B$ and

$$x \preceq_T^{\nu_0} y \implies \forall x' \forall y' (x' \preceq_T x \wedge y \preceq_T y' \implies x' \preceq_T^{\nu_0} y'). \quad (*)$$

Suppose that an ordinal $\nu_n \geq \nu_0$ satisfying $(*)$ has been defined. Put $C(x, y)$ iff $x \preceq_T^{\nu_n} y \wedge \forall z (y \preceq_T^{\nu_n} z \implies x \preceq_T^{\nu_n} z)$, so that C is a Π_1^1 set and $\preceq_T \subseteq C \subseteq \preceq_T^{\nu_n}$. As above there is an ordinal $\nu_{n+1} \geq \nu_n$ satisfying $\preceq_T \subseteq \preceq_T^{\nu_{n+1}} \subseteq C$, so that by definition

$$x \preceq_T^{\nu_{n+1}} y \implies \forall z (y \preceq_T^{\nu_n} z \implies x \preceq_T^{\nu_n} z).$$

Now take $\nu = \sup_n \nu_n$. □

Now assume $x \not\preceq_T y$ and prove that $\mathbf{h}(x) \neq \mathbf{h}(y)$. By the lemma there is an ordinal $\nu < \omega_1$ such that $x \not\preceq_T^\nu y$ and $\preceq_T^\nu$ is a p. q.-o.. It is a classical fact that the complement of $\preceq_T^\nu$ (which is the ν -th constituent of the co-analytic set $\mathbb{C}(\preceq_T)$) can be presented in the form of the set of all pairs $\langle x, y \rangle$ such that $C(x, y)$ is well-ordered and has the order type $\leq \nu$, where $C : \mathcal{N}^2 \longrightarrow \mathbb{Q}$ is a continuous function coded in $L[T] = L$. Therefore there exists a tree $S = S_\nu \in L$, $S \subseteq (\omega \times \omega \times \nu)^{<\omega}$, such that $\preceq_T^\nu = \mathbb{C}\mathbf{p}[S]$.

Apply Theorem 5 for the relations $\preceq_T = \mathbf{p}[T] \subseteq \preceq_T^\nu = \mathbb{C}\mathbf{p}[S]$. We observe that (II) of Theorem 5 fails by the assumption $(\dagger)$. Therefore (I) of Theorem 5 holds, so that there exists an ordinal $\alpha < \omega_1$ such that the map h_α satisfies $h_\alpha(x) = h_\alpha(y) \implies x \approx_T^\nu y$.

It follows that $h_\alpha(x) \neq h_\alpha(y)$ by the choice of ν , hence $\mathbf{h}(x) \neq \mathbf{h}(y)$, as required. Thus $\mathbf{h}$ witnesses (I^A) (the general case) of Theorem 2.

¹² Compare with the Burgess reflection theorem for Σ_1^1 equivalence relations.

8 Special cases: Borel classes and generic models

This section is devoted to the “additional” part in (I^A) of Theorem 2. *We continue to argue in the assumptions and notation of Subsection 7.2.* In particular we still assume (†) of Subsection 7.2, that is the given Σ_1^1 p. q.-o. $\preceq_{T=}\mathfrak{p}[T]$ does *not* satisfy (II^A) of Theorem 2.

We need here to specify definability properties related to the function **h**. An ordinary argument allows to define the sequence $\langle h_\alpha : \alpha < \omega_1 \rangle$ to be $\Delta_1^{L_{\omega_1}}$ hence Δ_1^{HC} . Then the map $\langle x, \gamma \rangle \mapsto \mathbf{h}(x) \upharpoonright \gamma$ is Δ_1^{HC} , too. Moreover there is a Σ_1 formula $\Phi(\cdot, \cdot, \cdot)$ such that

- (‡) If M is a transitive model of **ZFC**[−] (minus Power Sets), $x \in \mathcal{N} \cap M$, and $\lambda \in M$, $\lambda < \omega_1$, then $u = \mathbf{h}(x) \upharpoonright \lambda \in M$ and u is the only member of M such that $\Phi(x, \lambda, u)$ holds in M .

8.1 Order relations with Borel classes

Consider the case (I^A)(a) in the “additional” part of Theorem 2. Suppose that *every* $\approx_T$ -class $[x]_{\approx_T} = \{y : y \approx_T x\}$ is Borel. The aim is to find an antichain $A \subseteq 2^{<\omega_1}$ and a Δ_1^{HC} linearization $\mathbf{h}' : \langle \mathcal{N}; \preceq_T \rangle \longrightarrow \langle A; \leq_{\text{lex}} \rangle$.

First attempt. Let $x \in \mathcal{N}$. As $[x]_{\approx_T}$ is Borel, there exists, by Lemma 19, an ordinal $\lambda < \omega_1$ such that $x \approx_T y \iff x \approx_T^\lambda y$ and $\preceq_T^\lambda$ is a p. q.-o.. Therefore (see the end of Subsection 7.2) there is an ordinal $\alpha < \omega_1$ satisfying $h_\alpha(x) = h_\alpha(y) \implies x \approx_T^\nu y$, hence $h_\alpha(x) = h_\alpha(y) \iff x \approx_T y$ (for the chosen x and any y). It follows that $x \approx_T y \iff \mathbf{h}(x) \upharpoonright \lambda = \mathbf{h}(y) \upharpoonright \lambda$, for an ordinal $\lambda < \omega_1$. Let $\lambda_0(x)$ be the least such an ordinal and $\mathbf{h}_0(x) = \mathbf{h}(x) \upharpoonright \lambda_0(x)$. Now $A = \{\mathbf{h}_0(x) : x \in \mathcal{N}\}$ is an antichain in $2^{<\omega_1}$ while $\mathbf{h}_0 : \langle \mathcal{N}; \preceq_T \rangle \longrightarrow \langle A; \leq_{\text{lex}} \rangle$ is a linearization.

However the definition of $\lambda_0(x)$ seems not to provide that $\mathbf{h}_0$ is Δ_1^{HC} .

Second attempt. Let us modify the construction to meet the requirement that $\mathbf{h}_0$ is Δ_1^{HC} . The idea is to replace the quantifier $\forall y$ in the definition of $\lambda_0(x)$ by: *for each y in every set generic extension of a suitable model containing x* , which essentially means: *for comeager-many reals y* .

Let $x \in \mathcal{N}$. Define $\lambda_1(x)$ to be the least ordinal $\lambda < \omega_1$ such that

- (§) there exists a transitive model M of **ZFC**[−] containing λ , and a real $x' \in M$ satisfying $x \approx_T x'$ and, for any set generic extension M' of M and each real $z \in M'$, we have $\mathbf{h}(x') \upharpoonright \lambda = \mathbf{h}(z) \upharpoonright \lambda \implies x' \approx_T z$.

(Clearly $\lambda_1(x) \leq \lambda_0(x)$.)

Define $\mathbf{h}_1(x) = \mathbf{h}(x) \upharpoonright \lambda_1(x)$. Thus $\mathbf{h}_1 : \mathcal{N} \longrightarrow 2^{<\omega_1}$ is a Δ_1^{HC} function.

Lemma 20 *We have: $x \approx_T y$ iff $\mathbf{h}_1(x) = \mathbf{h}_1(y)$.*

Proof Suppose that $\mathbf{h}_1(x) = \mathbf{h}_1(y) = u \in 2^{<\omega_1}$ (the nontrivial direction), in particular $\lambda_1(x) = \lambda_1(y) = \lambda$. Let $x' \in M_x$ and $y' \in M_y$ witness that $\lambda_1(x) = \lambda_1(y) = \lambda$, in the sense of (§). Prove that $x' \approx_T y'$.

Let $f \in \lambda^\omega$ be a λ -collapse function generic over both M_x and M_y . Let ϑ_x, ϑ_y be the least ordinals not in resp. M_x, M_y . Suppose that $\vartheta_x \leq \vartheta_y$. Then both $M_x[f]$ and $L_{\vartheta_x}[f] \subseteq M_x[f]$ model $\mathbf{ZFC}^-$, hence by some version of Shoenfield there is a real $z \in L_{\vartheta_x}[f]$ satisfying $\mathbf{h}(z) \upharpoonright \lambda = u$, therefore $x' \approx_T z$ by the assumed (§). However z can be also considered in $L_{\vartheta_y}[f]$ which yields $y' \approx_T z$, hence $x' \approx_T y'$ as required. $\square$

Thus we have defined a Δ_1^{HC} map $\mathbf{h}_1 : \mathcal{N} \longrightarrow 2^{<\omega_1}$ such that, for any x , $\mathbf{h}_1(x) = \mathbf{h}(x) \upharpoonright \lambda_1(x)$ for some $\lambda_1(x) < \omega_1$, and $x \approx_T y \iff \mathbf{h}_1(x) = \mathbf{h}_1(y)$. However the range $A_1 = \{\mathbf{h}_1(x) : x \in \mathcal{N}\}$ may be not an antichain in $2^{<\omega_1}$.

To fix this last problem, we define a new Δ_1^{HC} map $\mathbf{h}' : \mathcal{N} \longrightarrow 5^{<\omega_1}$. First we change all values $\mathbf{h}_1(x)(\gamma) = 1$ to $\mathbf{h}'(x)(\gamma) = 4$ and add that $\mathbf{h}'(x)(\lambda_1(x)) = 2$. If $\nu < \lambda_1(x)$ is an ordinal of the form $\nu = \lambda_1(y)$ for some y then we change $\mathbf{h}'(x)(\nu)$ once again: if $x \preceq_T y$ then put $\mathbf{h}'(x)(\nu) = 1$ while if $y \preceq_T x$ then put $\mathbf{h}'(x)(\nu) = 3$. A simple verification shows that $\mathbf{h}'$ satisfies (I^A) of Theorem 2 (except for the fact that $\mathbf{h}'$ takes values in $5^{<\omega_1}$ rather than $2^{<\omega_1}$ but this can be easily fixed).

8.2 Order relations in a generic universe

This subsection is devoted to subitem (I^A)(b) in Theorem 2. In fact we shall assume the following: *there exists a real z_0 such that each real x in the universe V belongs to a set ¹³ generic extension of $L[z_0]$.* ¹⁴

It can be assumed that in fact $z_0 = 0$, so we simply drop z_0 .

Lemma 21 *Let $x \in \mathcal{N}$. There is an ordinal $\lambda < \omega_1$ such that*

- (i) $L_\lambda[\mathbf{h}(x) \upharpoonright \lambda]$ models $\mathbf{ZFC}^-$ (minus Power Sets);
- (ii) x belongs to a set generic extension of $L_\lambda[\mathbf{h}(x) \upharpoonright \lambda]$;
- (iii) if y, z are reals in a set generic extension M of $L_\lambda[\mathbf{h}(x) \upharpoonright \lambda]$, and $\mathbf{h}(y) \upharpoonright \lambda = \mathbf{h}(z) \upharpoonright \lambda$, then $y \approx_T z$.

¹³ It is not clear to what extent class forcing universes can accomodate the reasoning below, in particular the proofs of lemmas 21 and 22.

¹⁴ The extensions can be different for different reals x . Moreover the extensions can be Boolean valued extensions of $L[z]$ rather than factual classes in the universe.

Proof There exists a cardinal κ such that x belongs to a set generic extension of L_{κ^+} where κ^+ is taken in the sense of L . Note that L_{κ^+} models $\mathbf{ZFC}^-$. There is an ordinal $\lambda < \omega_1$ such that $\langle L_\lambda; x, \in \rangle$ is $\in$ -isomorphic to a countable elementary submodel of $\langle L_{\kappa^+}; x, \in \rangle$, hence L_λ models $\mathbf{ZFC}^-$ and x belongs to a set generic extension of L_λ . Note that $u = \mathbf{h}(x) \upharpoonright \lambda$ is a class in $L_\lambda[x]$ by $(\ddagger)$. Now it is a known fact (see Lemma 4.4 in Solovay [14] or Lemma 5 in Kanovei [10] as particular cases) that x belongs to a set generic extension of $L_\lambda[u]$.

Prove (iii). Suppose that reals y, z belong to a set generic extension of $L_\lambda[u]$ and $\mathbf{h}(y) \upharpoonright \lambda = \mathbf{h}(z) \upharpoonright \lambda$; prove $y \approx_T z$. First of all, a standard forcing argument shows that, as $u = \mathbf{h}(x) \upharpoonright \lambda$ is definable in $L_\lambda[x]$ while x belongs to a set generic extension of L_λ by the above, we can w. l. o. g. assume that y, z belong to a set generic extension of L_λ itself.

Let, for a transitive model M of $\mathbf{ZFC}^-$, $\mathbf{h}^M$ denote the map $\mathbf{h}$ defined in M (via e. g. the formula Φ of $(\ddagger)$). For instance $\mathbf{h}^V$ is simply $\mathbf{h}$. In any case $\mathbf{h}^M$ maps reals in M into $2^{\omega_1^M}$. In particular if each ordinal $\alpha \in M$ is countable in M and $\lambda = \text{Ord} \cap M$ then $\mathbf{h}^M(x) \in 2^\lambda$ for any real $x \in M$.

Note that (II^A) of Theorem 2 fails in any generic extension $L[G]$ of L because it is essentially a Σ_2^1 sentence (by Lemma 18) false in V by $(\dagger)$. Therefore, by the already proved, in Subsection 7.2, part of Theorem 2, every set forcing notion in L forces that, in the extension $L[G]$, for any two reals y, z , if $\mathbf{h}^{L[G]}(y) = \mathbf{h}^{L[G]}(z)$ then $y \approx_T z$.

A simple forcing argument transfers this result to L_{κ^+} as the initial model, and then, by the choice of λ , to L_λ , so that for any two reals y, z in a set generic extension $L_\lambda[G]$ of L_λ , if $\mathbf{h}^{L_\lambda[G]}(y) = \mathbf{h}^{L_\lambda[G]}(z)$ then $y \approx_T z$. It remains to note that $\mathbf{h}^{L_\lambda[G]}(y) = \mathbf{h}(y) \upharpoonright \omega_1^{L_\lambda[G]}$ for any real $y \in L_\lambda[G]$. $\square$

Let, for any $x \in \mathcal{N}$, λ_x be the least ordinal $\lambda < \omega_1$ satisfying the requirements of the lemma. We put $\mathbf{h}_1(x) = \mathbf{h}(x) \upharpoonright \lambda_x$. Apparently $\mathbf{h}_1$ is a Δ_1^{HC} map.

Lemma 22 *We have: $\mathbf{h}_1(x) = \mathbf{h}_1(y)$ iff $x \approx_T y$.*

Proof Suppose that $\mathbf{h}_1(x) = \mathbf{h}_1(y) = u \in 2^\lambda$ (so that $\lambda = \lambda_x = \lambda_y$) and prove $x \approx_T y$ (the nontrivial direction). Let x' and y' witness that $\mathbf{h}_1(x) = \mathbf{h}_1(y) = u$, so that they belong to resp. $L_\lambda[u, G_x]$ and $L_\lambda[u, G_y]$, which are resp. P_x -generic and P_y -generic extensions of $L_\lambda[u]$; P_x and P_y being set forcing notions in $L_\lambda[u]$. In addition, $x \approx_T x'$ and $y \approx_T y'$.

In particular, as $x \approx_T x'$, we have $\mathbf{h}(x) = \mathbf{h}(x')$, so that $u = \mathbf{h}(x') \upharpoonright \lambda$. Recall that the map $\mathbf{h}(\cdot) \upharpoonright \lambda$ results in some effective way from the $\Delta_1^{L_\lambda}$

sequence $\langle h_\alpha : \alpha < \lambda \rangle$. Therefore the fact that $\mathbf{h}(x') \restriction \lambda = u$ is forced in $L_\lambda[u]$. Thus we can assume that P_x forces in $L_\lambda[u]$ that $\mathbf{h}(x') \restriction \lambda = u$.

Consider a set $G \subseteq P_x$ which is P_x -generic over both $L_\lambda[u, G_x]$ and $L_\lambda[u, G_y]$. Let $z \in L_\lambda[u, G]$ be produced by G as x' from G_x , so that $\mathbf{h}(z) \restriction \lambda = u$. Thus two reals, x' and z , in the model $L_\lambda[u, G_x, G]$, satisfy $\mathbf{h}(z) \restriction \lambda = \mathbf{h}(x') \restriction \lambda$. It follows that $x' \approx_T z$ by the choice of λ .

We similarly prove $y' \approx_T z$, as required. $\square$

It remains to get $\mathbf{h}'$ from $\mathbf{h}_1$ as in Subsection 8.1.

$\square$ (*Theorem 2*)

References

- [1] S. D. Friedman and B. Velickovic. Nonstandard models and analytic equivalence relations. *Proc. Amer. Math. Soc.*, to appear.
- [2] L. A. Harrington, A. S. Kechris, A. Louveau. A Glimm – Effros dichotomy for Borel equivalence relations. *J. Amer. Math. Soc.* 1990, 3, pp. 903 – 928.
- [3] L. A. Harrington, D. Marker, S. Shelah, Borel orderings. *Trans. Amer. Math. Soc.* 1988, 310, pp. 293 – 302.
- [4] L. A. Harrington, S. Shelah, Counting equivalence classes for co- κ -Souslin equivalence relations. D. van Dalen *e. a.* eds, *Proceedings of the Conference in Prague* (LC'80), North Holland, 1982, pp. 147 – 152.
- [5] G. Hjorth. *A remark on Π_1^1 equivalence relations*. Note.
- [6] G. Hjorth. Thin equivalence relations and effective decompositions. *J. Symbolic Logic* 1993, 58, pp. 1153 – 1164.
- [7] G. Hjorth, A. S. Kechris. Analytic equivalence relations and Ulm-type classification. *J. Symbolic Logic* 1995, 60, pp. 1273 – 1299.
- [8] V. Kanovei. An Ulm-type classification theorem for equivalence relations in Solovay model. *J. Symbolic Logic* 1997, to appear.
- [9] V. Kanovei. Two dichotomy theorems on colourability of non-analytic graphs. *Fund. Math.*, to appear.
- [10] V. Kanovei. Ulm classification of analytic equivalence relations in generic universes. *Math. Logic Quarterly*, to appear.
- [11] V. Kanovei. When a partial Borel order is linearizable. *Fund. Math.*, submitted.

- [12] A. Louveau. Two results on Borel orders. *J. Symbolic Logic* 1989, 54, pp. 865 – 874.
- [13] S. Shelah. On co- κ -Souslin relations. *Israel J. Math* 1984, 47, pp. 139 – 153.
- [14] R. M. Solovay. A model of set theory in which every set of reals is Lebesgue measurable. *Ann. Math.* 1970, 92, pp. 1 – 56.